

FÖLNER TILINGS FOR ACTIONS OF AMENABLE GROUPS

CLINTON T. CONLEY, STEVE JACKSON, DAVID KERR, ANDREW MARKS,
BRANDON SEWARD, AND ROBIN TUCKER-DROB

ABSTRACT. We show that every probability-measure-preserving action of a countable amenable group G can be tiled, modulo a null set, using finitely many finite subsets of G (“shapes”) with prescribed approximate invariance so that the collection of tiling centers for each shape is Borel. This is a dynamical version of the Downarowicz–Huczek–Zhang tiling theorem for countable amenable groups and strengthens the Ornstein–Weiss Rokhlin lemma. As an application we prove that, for every countably infinite amenable group G , the crossed product of a generic free minimal action of G on the Cantor set is \mathcal{Z} -stable.

1. INTRODUCTION

A discrete group G is said to be *amenable* if it admits a finitely additive probability measure which is invariant under the action of G on itself by left translation, or equivalently if there exists a unital positive linear functional $\ell^\infty(G) \rightarrow \mathbb{C}$ which is invariant under the action of G on $\ell^\infty(G)$ induced by left translation (such a functional is called a *left invariant mean*). This definition was introduced by von Neumann in connection with the Banach–Tarski paradox and shown by Tarski to be equivalent to the absence of paradoxical decompositions of the group. Amenability has come to be most usefully leveraged through its combinatorial expression as the Følner property, which asks that for every finite set $K \subseteq G$ and $\delta > 0$ there exists a nonempty finite set $F \subseteq G$ which is (K, δ) -invariant in the sense that $|KF\Delta F| < \delta|F|$.

The concept of amenability appears as a common thread throughout much of ergodic theory as well as the related subject of operator algebras, where it is known via a number of avatars like injectivity, hyperfiniteness, and nuclearity. It forms the cornerstone of the theory of orbit equivalence, and also underpins both Kolmogorov–Sinai entropy and the classical ergodic theorems, whether explicitly in their most general formulations or implicitly in the original setting of single transformations (see Chapters 4 and 9 of [11]). A key tool in applying amenability to dynamics is the Rokhlin lemma of Ornstein and Weiss, which in one of its simpler forms says that for every free probability-measure-preserving action $G \curvearrowright (X, \mu)$ of a countably infinite amenable group and every finite set $K \subseteq G$ and $\delta > 0$ there exist (K, δ) -invariant finite sets $T_1, \dots, T_n \subseteq G$ and measurable sets $A_1, \dots, A_n \subseteq X$ such that the sets sA_i for $i = 1, \dots, n$ and $s \in T_i$ are pairwise disjoint and have union of measure at least $1 - \delta$ [17].

The proportionality in terms of which approximate invariance is expressed in the Følner condition makes it clear that amenability is a measure-theoretic property, and it is not

Date: March 27, 2017.

surprising that the most influential and definitive applications of these ideas in dynamics (e.g., the Connes–Feldman–Weiss theorem) occur in the presence of an invariant or quasi-invariant measure. Nevertheless, amenability also has significant ramifications for topological dynamics, for instance in guaranteeing the existence of invariant probability measures when the space is compact and in providing the basis for the theory of topological entropy. In the realm of operator algebras, similar comments can be made concerning the relative significance of amenability for von Neumann algebras (measure) and C^* -algebras (topology).

While the subjects of von Neumann algebras and C^* -algebras have long enjoyed a symbiotic relationship sustained in large part through the lens of analogy, and a similar relationship has historically bound together ergodic theory and topological dynamics, the last few years have witnessed the emergence of a new and structurally more direct kind of rapport between topology and measure in these domains, beginning on the operator algebra side with the groundbreaking work of Matui and Sato on strict comparison, \mathcal{Z} -stability, and decomposition rank [14, 15]. On the side of groups and dynamics, Downarowicz, Huczek, and Zhang recently showed that if G is a countable amenable group then for every finite set $K \subseteq G$ and $\delta > 0$ one can partition (or “tile”) G by left translates of finitely many (K, δ) -invariant finite sets [3]. The consequences that they derive from this tileability are topological and include the existence, for every such G , of a free minimal action with zero entropy. One of the aims of the present paper is to provide some insight into how these advances in operator algebras and dynamics, while seemingly unrelated at first glance, actually fit together as part of a common circle of ideas that we expect, among other things, to lead to further progress in the structure and classification theory of crossed product C^* -algebras.

Our main theorem is a version of the Downarowicz–Huczek–Zhang tiling result for free p.m.p. (probability-measure-preserving) actions of countable amenable groups which strengthens the Ornstein–Weiss Rokhlin lemma in the form recalled above by shrinking the leftover piece down to a null set (Theorem 3.6). As in the case of groups, one does not expect the utility of this dynamical tileability to be found in the measure setting, where the Ornstein–Weiss machinery generally suffices, but rather in the derivation of topological consequences. Indeed we will apply our tiling result to show that, for every countably infinite amenable group G , the crossed product $C(X) \rtimes G$ of a generic free minimal action $G \curvearrowright X$ on the Cantor set possesses the regularity property of \mathcal{Z} -stability (Theorem 5.4). The strategy is to first prove that such an action admits clopen tower decompositions with arbitrarily good Følner shapes (Theorem 4.2), and then to demonstrate that the existence of such tower decompositions implies that the crossed product is \mathcal{Z} -stable (Theorem 5.3). The significance of \mathcal{Z} -stability within the classification program for simple separable nuclear C^* -algebras is explained at the beginning of Section 5.

It is a curious irony in the theory of amenability that the Hall–Rado matching theorem can be used not only to show that the failure of the Følner property for a discrete group implies the formally stronger Tarski characterization of nonamenability in terms of the existence of paradoxical decompositions [2] but also to show, in the opposite direction, that the Følner property itself implies the formally stronger Downarowicz–Huczek–Zhang characterization of amenability which guarantees the existence of tilings of the group by translates of finitely many Følner sets [3]. This Janus-like scenario will be reprised

here in the dynamical context through the use of a measurable matching argument of Lyons and Nazarov that was originally developed to prove that for every simple bipartite nonamenable Cayley graph of a discrete group G there is a factor of a Bernoulli action of G which is an a.e. perfect matching of the graph [13]. Accordingly the basic scheme for proving Theorem 3.6 will be the same as that of Downarowicz, Huczek, and Zhang and divides into two parts:

- (i) using an Ornstein–Weiss-type argument to show that a subset of the space of lower Banach density close to one can be tiled by dynamical translates of Følner sets, and
- (ii) using a Lyons–Nazarov-type measurable matching to distribute almost all remaining points to existing tiles with only a small proportional increase in the size of the Følner sets, so that the approximate invariance is preserved.

We begin in Section 2 with the measurable matching result (Lemma 2.4), which is a variation on the Lyons–Nazarov theorem from [13] and is established along similar lines. In Section 3 we establish the appropriate variant of the Ornstein–Weiss Rokhlin lemma (Lemma 3.4) and put everything together in Theorem 3.6. Section 4 contains the genericity result for free minimal actions on the Cantor set, while Section 5 is devoted to the material on \mathcal{Z} -stability.

Acknowledgements. C.C. was partially supported by NSF grant DMS-1500906. D.K. was partially supported by NSF grant DMS-1500593. Part of this work was carried out while he was visiting the Erwin Schrödinger Institute (January–February 2016) and the Mittag-Leffler Institute (February–March 2016). A.M. was partially supported by NSF grant DMS-1500974. B.S. was partially supported by ERC grant 306494. R.T.D. was partially supported by NSF grant DMS-1600904. Part of this work was carried out during the AIM SQuaRE: Measurable Graph Theory.

2. MEASURABLE MATCHINGS

Given sets X and Y and a subset $\mathcal{R} \subseteq X \times Y$, with each $x \in X$ we associate its *vertical section* $\mathcal{R}_x = \{y \in Y : (x, y) \in \mathcal{R}\}$ and with each $y \in Y$ we associate its *horizontal section* $\mathcal{R}^y = \{x \in X : (x, y) \in \mathcal{R}\}$. Analogously, for $A \subseteq X$ we put $\mathcal{R}_A = \bigcup_{x \in A} \mathcal{R}_x = \{y \in Y : \exists x \in A (x, y) \in \mathcal{R}\}$. We say that \mathcal{R} is *locally finite* if for all $x \in X$ and $y \in Y$ the sets \mathcal{R}_x and \mathcal{R}^y are finite.

If now X and Y are standard Borel spaces equipped with respective Borel measures μ and ν , we say that $\mathcal{R} \subseteq X \times Y$ is *(μ, ν) -preserving* if whenever $f: A \rightarrow B$ is a Borel bijection between subsets $A \subseteq X$ and $B \subseteq Y$ with $\text{graph}(f) \subseteq \mathcal{R}$ we have $\mu(A) = \nu(B)$. We say that \mathcal{R} is *expansive* if there is some $c > 1$ such that for all Borel $A \subseteq X$ we have $\nu(\mathcal{R}_A) \geq c\mu(A)$.

We use the notation $f: X \rightarrow Y$ to denote a partial function from X to Y . We say that such a partial function f is *compatible* with $\mathcal{R} \subseteq X \times Y$ if $\text{graph}(f) \subseteq \mathcal{R}$.

Proposition 2.1 (ess. Lyons–Nazarov [13, Theorem 1.1]). *Suppose that X and Y are standard Borel spaces, that μ is a Borel probability measure on X , and that ν is a Borel measure on Y . Suppose that $\mathcal{R} \subseteq X \times Y$ is Borel, locally finite, (μ, ν) -preserving, and*

expansive. Then there is a μ -conull $X' \subseteq X$ and a Borel injection $f: X' \rightarrow Y$ compatible with \mathcal{R} .

Proof. Fix a constant of expansivity $c > 1$ for \mathcal{R} .

We construct a sequence $(f_n)_{n \in \mathbb{N}}$ of Borel partial injections from X to Y which are compatible with \mathcal{R} . Moreover, we will guarantee that the set $X' = \{x \in X : \exists m \in \mathbb{N} \forall n \geq m \ x \in \text{dom}(f_n) \text{ and } f_n(x) = f_m(x)\}$ is μ -conull, establishing that the limiting function satisfies the conclusion of the lemma.

Given a Borel partial injection $g: X \rightarrow Y$ we say that a sequence $(x_0, y_0, \dots, x_n, y_n) \in X \times Y \times \dots \times X \times Y$ is a g -augmenting path if

- $x_0 \in X$ is not in the domain of g ,
- for all distinct $i, j < n$, $y_i \neq y_j$.
- for all $i < n$, $y_i = g(x_{i+1})$,
- $y_n \in Y$ is not in the image of g .

We call n the *length* of such a g -augmenting path and x_0 the *origin* of the path. Note that the sequence $(x_0, y_0, y_1, \dots, y_n)$ in fact determines the entire g -augmenting path, and moreover that $x_i \neq x_j$ for distinct $i, j < n$.

In order to proceed we require the following two lemmas.

Lemma 2.2. *Suppose that $n \in \mathbb{N}$ and $g: X \rightarrow Y$ is a Borel partial injection compatible with \mathcal{R} admitting no augmenting paths of length less than n . Then $\mu(X \setminus \text{dom}(g)) \leq c^{-n}$.*

Proof. Put $A_0 = X \setminus \text{dom}(g)$. Define recursively for $i < n$ sets $B_i = \mathcal{R}_{A_i}$ and $A_{i+1} = A_i \cup g^{-1}(B_i)$. Note that the assumption that there are no augmenting paths of length less than n implies that each B_i is contained in the image of g . Expansivity of \mathcal{R} yields $\nu(B_i) \geq c\mu(A_i)$ and (μ, ν) -preservation of \mathcal{R} then implies that $\mu(A_{i+1}) \geq \nu(B_i) \geq c\mu(A_i)$. Consequently, $1 \geq \mu(A_n) \geq c^n \mu(A_0)$, and hence $\mu(A_0) \leq c^{-n}$. \square

Lemma 2.3 (ess. Elek-Lippner [5, Proposition 1.1]). *Suppose that $g: X \rightarrow Y$ is a Borel partial injection compatible with \mathcal{R} , and let $n \geq 1$. Then there is a Borel partial injection $g': X \rightarrow Y$ compatible with \mathcal{R} such that*

- $\text{dom}(g') \supseteq \text{dom}(g)$,
- g' admits no augmenting paths of length less than n ,
- $\mu(\{x \in X : g'(x) \neq g(x)\}) \leq n\mu(X \setminus \text{dom}(g))$.

Proof. Consider the set Z of injective sequences $(x_0, y_0, x_1, y_1, \dots, x_m, y_m)$, where $m < n$, such that for all $i \leq m$ we have $(x_i, y_i) \in \mathcal{R}$ and for all $i < m$ we have $(x_{i+1}, y_i) \in \mathcal{R}$. Equip Z with the standard Borel structure it inherits as a Borel subset of $(X \times Y)^{\leq n}$. Consider also the locally finite Borel graph \mathcal{G} on Z rendering adjacent two distinct sequences in Z if they share any entries. By [12, Proposition 4.5] there is a partition $Z = \bigsqcup_{k \in \mathbb{N}} Z_k$ of Z into Borel sets such that for all k , no two elements of Z_k are \mathcal{G} -adjacent. Fix a function $s: \mathbb{N} \rightarrow \mathbb{N}$ such that $s^{-1}(k)$ is infinite for all $k \in \mathbb{N}$.

Given a g -augmenting path $z = (x_0, y_0, \dots, x_m, y_m)$, define the *flip* along z to be the Borel partial function $g_z: X \rightarrow Y$ given by

$$g_z(x) = \begin{cases} y_i & \text{if } \exists i \leq m \ x = x_i, \\ g(x) & \text{otherwise.} \end{cases}$$

The fact that z is g -augmenting ensures that g_z is injective. More generally, for any Borel \mathcal{G} -independent set $Z^{\text{aug}} \subseteq Z$ of g -augmenting paths, we may simultaneously flip g along all paths in Z^{aug} to obtain another Borel partial injection $(g)_{Z^{\text{aug}}}$.

We iterate this construction. Put $g_0 = g$. Recursively assuming that $g_k: X \rightarrow Y$ has been defined, let Z_k^{aug} be the set of g_k -augmenting paths in $Z_{s(k)}$, and let $g_{k+1} = (g_k)_{Z_k^{\text{aug}}}$ be the result of flipping g_k along all paths in Z_k^{aug} . As each $x \in X$ is contained in only finitely many elements of Z , and since each path in Z can be flipped at most once (after the first flip its origin is always in the domain of the subsequent partial injections), it follows that the sequence $(g_k(x))_{k \in \mathbb{N}}$ is eventually constant. Defining $g'(x)$ to be the limiting value, it is routine to check that there are no g' -augmenting paths of length less than n .

Finally, to verify the third item of the lemma, put $A = \{x \in X : g'(x) \neq g(x)\}$. With each $x \in A$ associate the origin of the first augmenting path along which it was flipped. This is an at most n -to-1 Borel function from A to $X \setminus \text{dom}(g)$, and since \mathcal{R} is (μ, ν) -preserving the bound follows. \square

We are now in position to follow the strategy outlined at the beginning of the proof. Let $f_0: X \rightarrow Y$ be the empty function. Recursively assuming the Borel partial injection $f_n: X \rightarrow Y$ has been defined to have no augmenting paths of length less than n , let f_{n+1} be the Borel partial injection $(f_n)'$ granted by applying Lemma 2.3 to f_n . Thus f_{n+1} has no augmenting paths of length less than $n+1$ and the recursive construction continues.

Lemma 2.2 ensures that $\mu(X \setminus \text{dom}(f_n)) \leq c^{-n}$, and thus the third item of Lemma 2.3 ensures that $\mu(\{x \in X : f_{n+1}(x) \neq f_n(x)\}) \leq (n+1)c^{-n}$. As the sequence $(n+1)c^{-n}$ is summable, the Borel–Cantelli lemma implies that $X' = \{x \in X : \exists m \in \mathbb{N} \forall n \geq m \ x \in \text{dom}(f_n) \text{ and } f_n(x) = f_m(x)\}$ is μ -conull. Finally, $f = \lim_{n \rightarrow \infty} f_n \upharpoonright X'$ is as desired. \square

Lemma 2.4. *Suppose X and Y are standard Borel spaces, that μ is a Borel measure on X , and that ν is a Borel measure on Y . Suppose $\mathcal{R} \subseteq X \times Y$ is Borel, locally finite, (μ, ν) -preserving graph. Assume that there exist numbers $a, b > 0$ such that $|\mathcal{R}_x| \geq a$ for μ -a.e. $x \in X$ and $|\mathcal{R}^y| \leq b$ for ν -a.e. $y \in Y$. Then $\nu(\mathcal{R}_A) \geq \frac{a}{b}\mu(A)$ for all Borel subsets $A \subseteq X$.*

Proof. Since \mathcal{R} is (μ, ν) -preserving we have $\int_A |\mathcal{R}_x| d\mu = \int_{\mathcal{R}_A} |\mathcal{R}^y \cap A| d\nu$. Hence

$$a\mu(A) = \int_A a d\mu \leq \int_A |\mathcal{R}_x| d\mu = \int_{\mathcal{R}_A} |\mathcal{R}^y \cap A| d\nu \leq \int_{\mathcal{R}_A} b d\nu = b\nu(\mathcal{R}_A). \quad \square$$

3. FÖLNER TILINGS

Fix a countable group G . For finite sets $K, F \subseteq G$ and $\delta > 0$, we say that F is (K, δ) -invariant if $|KF\Delta F| < \delta|F|$. Note this condition implies $|KF| < (1 + \delta)|F|$. Recall that G is *amenable* if for every finite $K \subseteq G$ and $\delta > 0$ there exists a (K, δ) -invariant set F . A *Følner sequence* is a sequence of finite sets $F_n \subseteq G$ with the property that for every finite $K \subseteq G$ and $\delta > 0$ the set F_n is (K, δ) -invariant for all but finitely many n . Below, we always assume that G is amenable.

Fix a free action $G \curvearrowright X$. For $A \subseteq X$ we define the lower and upper *Banach densities* of A to be

$$\underline{D}(A) = \sup_{\substack{F \subseteq G \\ F \text{ finite}}} \inf_{x \in X} \frac{|A \cap Fx|}{|F|} \quad \text{and} \quad \bar{D}(A) = \inf_{\substack{F \subseteq G \\ F \text{ finite}}} \sup_{x \in X} \frac{|A \cap Fx|}{|F|}.$$

Equivalently [3, Lem. 2.9], if $(F_n)_{n \in \mathbb{N}}$ is a Følner sequence then

$$\underline{D}(A) = \lim_{n \rightarrow \infty} \inf_{x \in X} \frac{|A \cap F_n x|}{|F_n|} \quad \text{and} \quad \bar{D}(A) = \lim_{n \rightarrow \infty} \sup_{x \in X} \frac{|A \cap F_n x|}{|F_n|}.$$

We now define an analogue of ‘ (K, δ) -invariant’ for infinite subsets of X . A set $A \subseteq X$ (possibly infinite) is $(K, \delta)^*$ -invariant if there is a finite set $F \subseteq G$ such that $|(KA\Delta A) \cap Fx| < \delta|A \cap Fx|$ for all $x \in X$. Equivalently, A is $(K, \delta)^*$ -invariant if and only if for every Følner sequence $(F_n)_{n \in \mathbb{N}}$ we have $\lim_n \sup_x |(KA\Delta A) \cap F_n x|/|A \cap F_n x| < \delta$.

A collection $\{F_n : n \in \mathbb{N}\}$ of finite subsets of X is called ϵ -disjoint if for each n there is $F'_n \subseteq F_n$ such that $|F'_n| > (1 - \epsilon)|F_n|$ and such that the sets $\{F'_n : n \in \mathbb{N}\}$ are pairwise disjoint.

Lemma 3.1. *Let $K, W \subseteq G$ be finite, let $\epsilon, \delta > 0$, let $C \subseteq X$, and for $c \in C$ let $F_c \subseteq W$ be $(K, \delta(1 - \epsilon))$ -invariant. If the collection $\{F_c c : c \in C\}$ is ϵ -disjoint and $\bigcup_{c \in C} F_c c$ has positive lower Banach density, then $\bigcup_{c \in C} F_c c$ is $(K, \delta)^*$ -invariant.*

Proof. Set $A = \bigcup_{c \in C} F_c c$ and set $T = WW^{-1}(\{1_G\} \cup K)^{-1}$. Since W is finite and each $F_c \subseteq W$, there is $0 < \delta_0 < \delta$ such that each F_c is $(K, \delta_0(1 - \epsilon))$ -invariant. Fix a finite set $U \subseteq G$ which is $(T, \frac{D(A)}{2|T|}(\delta - \delta_0))$ -invariant and satisfies $\inf_{x \in X} |A \cap Ux| > \frac{D(A)}{2}|U|$. Now fix $x \in X$. Let B be the set of $b \in Ux$ such that $Tb \not\subseteq Ux$. Note that $B \subseteq T^{-1}(TU\Delta Ux)$ and thus

$$|B| \leq |T| \cdot \frac{|TU\Delta U|}{|U|} \cdot \frac{|U|}{|A \cap Ux|} \cdot |A \cap Ux| < (\delta - \delta_0)|A \cap Ux|.$$

Set $C' = \{c \in C : F_c c \subseteq Ux\}$. Note that the ϵ -disjoint assumption gives $(1 - \epsilon) \sum_{c \in C'} |F_c| \leq |A \cap Ux|$. Also, our definitions of C' , T , and B imply that if $c \in C \setminus C'$ and $(\{1_G\} \cup K)F_c c \cap Ux \neq \emptyset$ then $((\{1_G\} \cup K)F_c c) \cap Ux \subseteq B$. Therefore $(KA\Delta A) \cap Ux \subseteq B \cup \bigcup_{c \in C'} (KF_c c \Delta F_c c)$. Combining this with the fact that each set F_c is $(K, \delta_0(1 - \epsilon))$ -invariant, we obtain

$$\begin{aligned} |(KA\Delta A) \cap Ux| &\leq |B| + \sum_{c \in C'} |KF_c c \Delta F_c c| \\ &< (\delta - \delta_0)|A \cap Ux| + \sum_{c \in C'} \delta_0(1 - \epsilon)|F_c| \\ &\leq (\delta - \delta_0)|A \cap Ux| + \delta_0|A \cap Ux| \\ &= \delta|A \cap Ux|. \end{aligned}$$

Since x was arbitrary, we conclude that A is $(K, \delta)^*$ -invariant. \square

Lemma 3.2. *Let $T \subseteq G$ be finite and let $\epsilon, \delta > 0$ with $\epsilon(1 + \delta) < 1$. Suppose that $A \subseteq X$ is $(T^{-1}, \delta)^*$ -invariant. If $B \supseteq A$ and $|B \cap Tx| \geq \epsilon|T|$ for all $x \in X$, then*

$$\underline{D}(B) \geq (1 - \epsilon(1 + \delta)) \cdot \underline{D}(A) + \epsilon.$$

Proof. This is implicitly demonstrated in [3, Proof of Lem. 4.1]. As a convenience to the reader, we include a proof here. Fix $\theta > 0$. Since A is $(T^{-1}, \delta)^*$ -invariant, we can pick a finite set $U \subseteq G$ which is (T, θ) -invariant and satisfies

$$\inf_{x \in X} \frac{|A \cap Ux|}{|U|} > D(A) - \theta \quad \text{and} \quad \sup_{x \in X} \frac{|T^{-1}A \cap Ux|}{|A \cap Ux|} < 1 + \delta.$$

Fix $x \in X$, set $\alpha = \frac{|A \cap Ux|}{|U|} > D(A) - \theta$, and set $U' = \{u \in U : A \cap Tux = \emptyset\}$. Notice that

$$\frac{|U'|}{|U|} = \frac{|U| - |T^{-1}A \cap Ux|}{|U|} = 1 - \frac{|T^{-1}A \cap Ux|}{|A \cap Ux|} \cdot \frac{|A \cap Ux|}{|U|} > 1 - (1 + \delta)\alpha.$$

Since $A \cap TU'x = \emptyset$ and $|B \cap Ty| \geq \epsilon|T|$ for all $y \in X$, it follows that $|(B \setminus A) \cap Tux| \geq \epsilon|T|$ for all $u \in U'$. Thus there are $\epsilon|T||U'|$ many pairs $(t, u) \in T \times U'$ with $tux \in B \setminus A$. It follows there is $t^* \in T$ with $|(B \setminus A) \cap t^*U'x| \geq \epsilon \cdot |U'|$. Therefore

$$\begin{aligned} \frac{|B \cap TUX|}{|TU|} &\geq \left(\frac{|A \cap Ux|}{|U|} + \frac{|(B \setminus A) \cap t^*U'x|}{|U'|} \cdot \frac{|U'|}{|U|} \right) \cdot \frac{|U|}{|TU|} \\ &> \left(\alpha + \epsilon(1 - (1 + \delta)\alpha) \right) \cdot (1 + \theta)^{-1} \\ &= \left((1 - \epsilon(1 + \delta))\alpha + \epsilon \right) \cdot (1 + \theta)^{-1} \\ &> \left((1 - \epsilon(1 + \delta))(D(A) - \theta) + \epsilon \right) \cdot (1 + \theta)^{-1}. \end{aligned}$$

Letting θ tend to 0 completes the proof. \square

Lemma 3.3. *Let X be a standard Borel space and let $G \curvearrowright X$ be a free Borel action. Let $Y \subseteq X$ be Borel, let $T \subseteq G$ be finite, and let $\epsilon \in (0, 1/2)$. Then there is a Borel set $C \subseteq X$ and a Borel function $c \in C \mapsto T_c \subseteq T$ such that $|T_c| > (1 - \epsilon)|T|$, the sets $\{T_{c_c} : c \in C\}$ are pairwise disjoint and disjoint with Y , $Y \cup \bigcup_{c \in C} T_c c = Y \cup TC$, and $|(Y \cup TC) \cap Tx| \geq \epsilon|T|$ for all $x \in X$.*

Proof. Fix a Borel partition $\mathcal{P} = \{P_1, \dots, P_m\}$ of X such that $Tx \cap Tx' = \emptyset$ for all $x \neq x' \in P_i$ and all $1 \leq i \leq m$. We will pick Borel sets $C_i \subseteq P_i$ and set $C = \bigcup_{1 \leq i \leq m} C_i$. Set $Y_0 = Y$. Let $1 \leq i \leq m$ and inductively assume that Y_{i-1} has been defined. Define $C_i = \{c \in P_i : |Y_{i-1} \cap Tc| < \epsilon|T|\}$, define $Y_i = Y_{i-1} \cup TC_i$, and for $c \in C_i$ set $T_c = \{t \in T : tc \notin Y_{i-1}\}$. It is easily seen that $C = \bigcup_{1 \leq i \leq m} C_i$ has the desired properties. \square

The following lemma is mainly due to Ornstein–Weiss [17], who proved this with an invariant probability measure taking the place of Banach density. It was adapted to the Banach density setting by Downarowicz–Huczuk–Zhang [3]. The only difference between this lemma and prior versions is we work both in the Borel setting and use Banach density.

Lemma 3.4. [17, I.§2. Thm. 6] [3, Lem. 4.1] *Let X be a standard Borel space and let $G \curvearrowright X$ be a free Borel action. Let $K \subseteq G$ be finite, let $\epsilon \in (0, 1/2)$, and let n satisfy $(1 - \epsilon)^n < \epsilon$. Then there exist (K, ϵ) -invariant sets F_1, \dots, F_n , a Borel set $C \subseteq X$, and a Borel function $c \in C \mapsto F_c \subseteq G$ such that:*

- (i) *for every $c \in C$ there is $1 \leq i \leq n$ with $F_c \subseteq F_i$ and $|F_c| > (1 - \epsilon)|F_i|$;*
- (ii) *the sets F_{c_c} , $c \in C$, are pairwise disjoint; and*

(iii) $D(\bigcup_{c \in C} F_c c) > 1 - \epsilon$.

Proof. Fix $\delta > 0$ satisfying $(1 + \delta)^{-1}(1 - (1 + \delta)\epsilon)^n < \epsilon - 1 + (1 + \delta)^{-1}$. Fix a sequence of (K, ϵ) -invariant sets F_1, \dots, F_n such that F_i is $(F_j^{-1}, \delta(1 - \epsilon))$ -invariant for all $1 \leq j < i \leq n$.

The set C will be the disjoint union of sets C_i , $1 \leq i \leq n$. The construction will be such that $F_c \subseteq F_i$ and $|F_c| > (1 - \epsilon)|F_i|$ for $c \in C_i$. We will define $A_i = \bigcup_{i \leq k \leq n} \bigcup_{c \in C_k} F_c c$ and arrange that $A_{i+1} \cup F_i C_i = A_{i+1} \cup \bigcup_{c \in C_i} F_c c$ and

$$(3.1) \quad D(A_i) \geq (1 + \delta)^{-1} - (1 + \delta)^{-1}(1 - \epsilon(1 + \delta))^{n+1-i}.$$

In particular, we will have $A_i = \bigcup_{i \leq k \leq n} F_k C_k$.

To begin, apply Lemma 3.3 with $Y = \emptyset$ and $T = F_n$ to get a Borel set C_n and a Borel map $c \in C_n \mapsto F_c \subseteq F_n$ such that $|F_c| > (1 - \epsilon)|F_n|$, the sets $\{F_c c : c \in C_n\}$ are pairwise disjoint, $\bigcup_{c \in C_n} F_c c = F_n C_n$, and $|F_n C_n \cap F_n x| \geq \epsilon|F_n|$ for all $x \in X$. Applying Lemma 3.2 with $A = \emptyset$ and $B = F_n C_n$ we find that the set $A_n = F_n C_n$ satisfies $D(A_n) \geq \epsilon$.

Inductively assume that C_n through C_{i+1} have been defined and A_n through A_{i+1} are defined as above and satisfy (3.1). Using $Y = A_{i+1}$ and $T = F_i$, apply Lemma 3.3 to get a Borel set C_i and a Borel map $c \in C_i \mapsto F_c \subseteq F_i$ such that $|F_c| > (1 - \epsilon)|F_i|$, the sets $\{F_c c : c \in C_i\}$ are pairwise disjoint and disjoint with A_{i+1} , $A_{i+1} \cup \bigcup_{c \in C_i} F_c c = A_{i+1} \cup F_i C_i$, and $|(A_{i+1} \cup F_i C_i) \cap F_i x| \geq \epsilon|F_i|$ for all $x \in X$. The set A_{i+1} is the union of an ϵ -disjoint collection of $(F_i^{-1}, \delta(1 - \epsilon))$ -invariant sets and has positive lower Banach density. So by Lemma 3.1 A_{i+1} is $(F_i^{-1}, \delta)^*$ -invariant. Applying Lemma 3.2 with $A = A_{i+1}$, we find that $A_i = A_{i+1} \cup F_i C_i$ satisfies

$$\begin{aligned} D(A_i) &\geq (1 - \epsilon(1 + \delta)) \cdot D(A_{i+1}) + \epsilon \\ &\geq \frac{(1 - \epsilon(1 + \delta))}{(1 + \delta)} - (1 + \delta)^{-1}(1 - \epsilon(1 + \delta))^{n+1-i} + \frac{\epsilon(1 + \delta)}{1 + \delta} \\ &= (1 + \delta)^{-1} - (1 + \delta)^{-1}(1 - \epsilon(1 + \delta))^{n+1-i}. \end{aligned}$$

This completes the inductive step and completes the definition of C . It is immediate from the construction that (i) and (ii) are satisfied. Clause (iii) also follows by noting that (3.1) is greater than $1 - \epsilon$ when $i = 1$. \square

We recall the following simple fact.

Lemma 3.5. [3, Lem. 2.3] *If $F \subseteq G$ is (K, δ) -invariant and F' satisfies $|F' \triangle F| < \epsilon|F|$ then F' is $(K, \frac{(|K|+1)\epsilon+\delta}{1-\epsilon})$ -invariant.*

Now we present the main theorem.

Theorem 3.6. *Let G be a countable amenable group, let (X, μ) be a standard probability space, and let $G \curvearrowright (X, \mu)$ be a free p.m.p. action. For every finite $K \subseteq G$ and every $\delta > 0$ there exist a μ -conull G -invariant Borel set $X' \subseteq X$, a collection $\{C_i : 0 \leq i \leq m\}$ of Borel subsets of X' , and a collection $\{F_i : 0 \leq i \leq m\}$ of (K, δ) -invariant sets such that $\{F_i c : 0 \leq i \leq m, c \in C_i\}$ partitions X' .*

Proof. Fix $\epsilon \in (0, 1/2)$ satisfying $\frac{(|K|+1)6\epsilon+\epsilon}{1-6\epsilon} < \delta$. Apply Lemma 3.4 to get (K, ϵ) -invariant sets F'_1, \dots, F'_n , a Borel set $C \subseteq X$, and a Borel function $c \in C \mapsto F_c \subseteq G$ satisfying

- (i) for every $c \in C$ there is $1 \leq i \leq n$ with $F_c \subseteq F'_i$ and $|F_c| > (1 - \epsilon)|F'_i|$;
- (ii) the sets $F_c c$, $c \in C$, are pairwise disjoint; and
- (iii) $D(\bigcup_{c \in C} F_c c) > 1 - \epsilon$.

Set $Y = X \setminus \bigcup_{c \in C} F_c c$. If $\mu(Y) = 0$ then we are done. So we assume $\mu(Y) > 0$ and we let ν denote the restriction of μ to Y . Fix a Borel map $c \in C \mapsto Z_c \subseteq F_c$ satisfying $4\epsilon|F_c| < |Z_c| < 5\epsilon|F_c|$ for all $c \in C$ (its clear from the proof of Lemma 3.4 that we may choose the sets F'_i so that $\epsilon|F_c| > \min_i \epsilon(1 - \epsilon)|F'_i| > 1$). Set $Z = \bigcup_{c \in C} Z_c c$ and let ζ denote the restriction of μ to Z (note that $\mu(Z) > 0$).

Set $W = \bigcup_{i=1}^n F'_i$ and $W' = WW^{-1}$. Fix a finite set $U \subseteq G$ which is $(W', (1/2 - \epsilon)/|W'|)$ -invariant and satisfies $\inf_{x \in X} |(X \setminus Y) \cap Ux| > (1 - \epsilon)|U|$. Since every amenable group admits a Følner sequence consisting of symmetric sets, we may assume that $U = U^{-1}$ [16, Cor. 5.3]. Define $\mathcal{R} \subseteq Y \times Z$ by declaring $(y, z) \in \mathcal{R}$ if and only if $y \in Uz$ (equivalently $z \in Uy$). Then \mathcal{R} is Borel, locally finite, and (ν, ζ) -preserving. We now check that \mathcal{R} is expansive. We automatically have $|\mathcal{R}^z| = |Y \cap Uz| < \epsilon|U|$ for all $z \in Z$. By Lemma 2.4 it suffices to show that $|\mathcal{R}_y| = |Z \cap Uy| \geq 2\epsilon|U|$ for all $y \in Y$. Fix $y \in Y$. Let B be the set of $b \in Uy$ such that $W'b \not\subseteq Uy$. Then $B \subseteq W'(W'Uy \Delta Uy)$ and thus

$$\frac{|B|}{|U|} \leq |W'| \cdot \frac{|W'U \Delta U|}{|U|} < 1/2 - \epsilon.$$

Let A be the union of those sets $F_c c$, $c \in C$, which are contained in Uy . Notice that $(X \setminus Y) \cap Uy \subseteq B \cup A$. Therefore

$$\frac{1}{2} - \epsilon + \frac{|A|}{|U|} > \frac{|(B \cup A) \cap Uy|}{|U|} \geq \frac{|(X \setminus Y) \cap Uy|}{|U|} > 1 - \epsilon,$$

hence $|A| > |U|/2$. By construction $|Z \cap A| > 4\epsilon|A|$. So $|Z \cap Uy| \geq |Z \cap A| > 2\epsilon|U|$. We conclude that \mathcal{R} is expansive.

Apply Proposition 2.1 to obtain a G -invariant μ -conull set $X' \subseteq X$ and a Borel injection $\rho : Y \cap X' \rightarrow Z$ with $\text{graph}(\rho) \subseteq \mathcal{R}$. Consider the sets $F_c \cup \{g \in U : gc \in Y \text{ and } \rho(gc) \in F_c c\}$ as $c \in C$ varies. These are subsets of $W \cup U$ and thus there are only finitely many such sets which we can enumerate as F_1, \dots, F_m . We partition $C \cap X'$ into Borel sets C_1, \dots, C_m with $c \in C_i$ if and only if $c \in X'$ and $F_c \cup \{g \in U : gc \in Y \text{ and } \rho(gc) \in F_c c\} = F_i$. Since ρ is defined on all of $Y \cap X'$, we see that the sets $\{F_i c : 1 \leq i \leq m, c \in C_i\}$ partition X' . Finally, for $c \in C_i \cap X'$, if we let F'_j be such that $|F_c \Delta F'_j| < \epsilon|F'_j|$, then

$$\begin{aligned} |F_i \Delta F'_j| &\leq |F_i \Delta F_c| + |F_c \Delta F'_j| \leq |\rho^{-1}(F_c c)| + \epsilon|F'_j| \\ &\leq |Z_c| + \epsilon|F'_j| < 5\epsilon|F_c| + \epsilon|F'_j| \leq 6\epsilon|F'_j|. \end{aligned}$$

Using Lemma 3.5 and our choice of ϵ , this implies that each set F_i is (K, δ) -invariant. \square

4. CLOPEN TOWER DECOMPOSITIONS WITH FÖLNER SHAPES

Let $G \curvearrowright X$ be an action of a group on a compact space. By a *clopen tower* we mean a pair (B, S) where B is a clopen subset of X (the *base* of the tower) and S is a finite subset of G (the *shape* of the tower) such that the sets sB for $s \in S$ are pairwise disjoint. By a *clopen tower decomposition* of X we mean a finite collection $\{(B_i, S_i)\}_{i=1}^n$ of clopen towers such that the sets $S_1 B_1, \dots, S_n B_n$ form a partition of X . We also similarly speak of *measurable towers* and *measurable tower decompositions* for an action $G \curvearrowright (X, \mu)$ on a

measure space, with the bases now being measurable sets instead of clopen sets. In this terminology, Theorem 3.6 says that if $G \curvearrowright (X, \mu)$ is a free p.m.p. action of a countable amenable group on a standard probability space then for every finite set $K \subseteq G$ and $\delta > 0$ there exists, modulo a null set, a measurable tower decomposition of X with (K, δ) -invariant shapes.

Lemma 4.1. *Let G be a countably infinite amenable group and $G \curvearrowright X$ a free minimal action on the Cantor set. Then this action has a free minimal extension $G \curvearrowright Y$ on the Cantor set such that for every finite set $F \subseteq G$ and $\delta > 0$ there is a clopen tower decomposition of Y with (F, δ) -invariant shapes.*

Proof. Let $F_1 \subseteq F_2 \subseteq \dots$ be an increasing sequence of finite subsets of G whose union is equal to G . Fix a G -invariant Borel probability measure μ on X (such a measure exists by amenability). The freeness of the action $G \curvearrowright X$ means that for each $n \in \mathbb{N}$ we can apply Theorem 3.6 to produce, modulo a null set, a measurable tower decomposition \mathcal{U}_n for the p.m.p. action $G \curvearrowright (X, \mu)$ such that each shape is $(F_n, 1/n)$ -invariant. Let A be the unital G -invariant C^* -algebra of $L^\infty(X, \mu)$ generated by $C(X)$ and the indicator functions of the levels of each of the tower decompositions \mathcal{U}_n . Since there are countably many such indicator functions and the group G is countable, the C^* -algebra A is separable. Therefore by the Gelfand-Naimark theorem we have $A = C(Z)$ for some zero-dimensional metrizable space Z and a G -factor map $\varphi : Z \rightarrow X$. By a standard fact which can be established using Zorn's lemma, there exists a nonempty closed G -invariant set $Y \subseteq Z$ such that the restriction action $G \curvearrowright Y$ is minimal. Note that Y is necessarily a Cantor set, since G is infinite. Also, the action $G \curvearrowright Y$ is free, since it is an extension of a free action. Since the action on Y is minimal, the restriction $\varphi|_Y : Y \rightarrow X$ is surjective and hence a G -factor map. For each n we get from \mathcal{U}_n a clopen tower decomposition \mathcal{V}_n of Y with $(F_n, 1/n)$ -invariant shapes, and by intersecting the levels of the towers in \mathcal{V}_n with Y we obtain a clopen tower decomposition of Y with $(F_n, 1/n)$ -invariant shapes, showing that the extension $G \curvearrowright Y$ has the desired property. \square

Let X be the Cantor set and let G be a countable infinite amenable group. The set $\text{Act}(G, X)$ is a Polish space under the topology which has as a basis the sets

$$U_{\alpha, \mathcal{P}, F} = \{\beta \in \text{Act}(G, X) : \alpha_s A = \beta_s A \text{ for all } A \in \mathcal{P} \text{ and } s \in F\}$$

where $\alpha \in \text{Act}(G, X)$, \mathcal{P} is a clopen partition of X , and F is a finite subset of G . Write $\text{FrMin}(G, X)$ for the set of actions in $\text{Act}(G, X)$ which are free and minimal. Then $\text{FrMin}(G, X)$ is a G_δ set. To see this, fix a sequence $F_1 \subseteq F_2 \subseteq \dots$ of finite subsets of G with union equal to G and an enumeration s_1, s_2, s_3, \dots of G , and for every $n \in \mathbb{N}$ and nonempty clopen set $A \subseteq X$ define the set $\mathcal{W}_{n,A}$ of all $\alpha \in \text{Act}(G, X)$ such that (i) $\bigcup_{s \in F_n} \alpha_s A = X$ and (ii) there exists a clopen partition $\{A_1, \dots, A_k\}$ of A such that $\alpha_{s_n} A_i \cap A_i = \emptyset$ for all $i = 1, \dots, k$. Then each $\mathcal{W}_{n,A}$ is open, which means, with A ranging over the countable collection of nonempty clopen subsets of X , that the intersection $\bigcap_{n \in \mathbb{N}} \bigcap_A \mathcal{W}_{n,A}$, which is equal to $\text{FrMin}(G, X)$, is a G_δ set. It follows that $\text{FrMin}(G, X)$ is a Polish space.

Theorem 4.2. *Let G be a countably infinite amenable group. Let \mathcal{C} be the collection of actions in $\text{FrMin}(G, X)$ with the property that for every finite set $F \subseteq G$ and $\delta > 0$ there*

is a clopen tower decomposition of X with (F, δ) -invariant shapes. Then \mathcal{C} is a dense G_δ subset of $\text{FrMin}(G, X)$.

Proof. That \mathcal{C} is a G_δ set is a simple exercise. Let $G \curvearrowright^\alpha X$ be any action in $\text{FrMin}(G, X)$. By Lemma 4.1 this action has a free minimal extension $G \curvearrowright^\beta Y$ with the property in the theorem statement, where Y is the Cantor set. Let \mathcal{P} be a clopen partition of X and F a nonempty finite subset of G . Write A_1, \dots, A_n for the members of the clopen partition $\bigvee_{s \in F} s^{-1}\mathcal{P}$. Then for each $i = 1, \dots, n$ the set A_i and its inverse image $\varphi^{-1}(A_i)$ under the extension map $\varphi : Y \rightarrow X$ are Cantor sets, and so we can find a homeomorphism $\psi_i : A_i \rightarrow \varphi^{-1}(A_i)$. Let $\psi : X \rightarrow Y$ be the homeomorphism which is equal to ψ_i on A_i for each i . Then the action $G \curvearrowright^\gamma X$ defined by $\gamma_s = \psi^{-1} \circ \beta_s \circ \psi$ for $s \in G$ belongs to \mathcal{C} as well as to the basic open neighborhood $U_{\alpha, \mathcal{P}, F}$ of α , establishing the density of \mathcal{C} . \square

5. APPLICATIONS TO \mathcal{Z} -STABILITY

A C^* -algebra A is said to be \mathcal{Z} -stable if $A \otimes \mathcal{Z} \cong A$ where \mathcal{Z} is the Jiang–Su algebra [10], with the C^* -tensor product being unique in this case because \mathcal{Z} is nuclear. \mathcal{Z} -stability has become an important regularity property in the classification program for simple separable nuclear C^* -algebras, which has recently witnessed some spectacular advances. Thanks to recent work of Elliott–Gong–Lin–Niu [6, 4] and Tikuisis–White–Winter [21], it is now known that simple separable unital C^* -algebras satisfying the universal coefficient theorem and having finite nuclear dimension are classified by ordered K -theory paired with tracial states. Although \mathcal{Z} -stability does not appear in the hypotheses of this classification theorem, it does play an important technical role in the proof. Moreover, it is a conjecture of Toms and Winter that for simple separable infinite-dimensional unital nuclear C^* -algebras the following properties are equivalent:

- (i) \mathcal{Z} -stability,
- (ii) finite nuclear dimension,
- (iii) strict comparison.

Implications between (i), (ii), and (iii) are known to hold in various degrees of generality. In particular, the implication (ii) \Rightarrow (i) was established in [22] while the converse is known to hold when the extreme boundary of the convex set of tracial states is compact [1]. It remains a problem to determine whether any of the crossed products of the actions in Theorem 5.4 falls within the purview of these positive results on the Toms–Winter conjecture, and in particular whether any of them has finite nuclear dimension (see Question 5.5).

By now there exist highly effectively methods for establishing finite nuclear dimension for crossed products of free actions on compact metrizable spaces of finite covering dimension [19, 20, 7], but their utility is structurally restricted to groups with finite asymptotic dimension and hence excludes many amenable examples like the Grigorchuk group. One can show using the technology from [7] that, for a countably infinite amenable group with finite asymptotic dimension, the crossed product of a generic free minimal action on the Cantor set has finite nuclear dimension. Our intention here has been to remove the restriction of finite asymptotic dimension by means of a different approach that establishes instead the conjecturally equivalent property of \mathcal{Z} -stability but for arbitrary countably infinite amenable groups.

To simplify the verification of \mathcal{Z} -stability in the proof of Theorem 5.3 we will use the following result from [8], which is based on methods from [14]. We state it here with stronger but simpler hypotheses that are sufficient for our purposes.

Theorem 5.1. *Let A be a simple separable unital nuclear C^* -algebra. Suppose that for every $n \in \mathbb{N}$, finite set $\Omega \subseteq A$, and $\varepsilon > 0$ there exist an order-zero complete positive contractive linear map $\varphi : M_n \rightarrow A$ and a $v \in A$ such that*

- (i) $vv^* = 1 - \varphi(1)$,
- (ii) $v^*v \leq \varphi(e_{11})$,
- (iii) $\|[a, \varphi(b)]\| < \varepsilon$ for all $a \in \Omega$ and norm-one $b \in M_n$.

Then A is \mathcal{Z} -stable.

The following is the Ornstein–Weiss quasitiling theorem [17] as formulated in Theorem 3.36 of [11].

Theorem 5.2. *Let $0 < \beta < \frac{1}{2}$ and let n be a positive integer such that $(1 - \beta/2)^n < \beta$. Then whenever $e \in T_1 \subseteq T_2 \subseteq \dots \subseteq T_n$ are finite subsets of a group G such that $|\partial_{T_{i-1}}T_i| \leq (\eta/8)|T_i|$ for $i = 2, \dots, n$, for every $(T_n, \beta/4)$ -invariant nonempty finite set $E \subseteq G$ there exist $C_1, \dots, C_n \subseteq G$ such that*

- (i) $\bigcup_{i=1}^n T_i C_i \subseteq E$, and
- (ii) the collection of right translates $\bigcup_{i=1}^n \{T_i c : c \in C_i\}$ is β -disjoint and $(1 - \beta)$ -covers E .

Theorem 5.3. *Let G be a countably infinite amenable group and let $G \curvearrowright X$ be a free minimal action on the Cantor set such that for every finite set $F \subseteq G$ and $\delta > 0$ there is a clopen tower decomposition of X with (F, δ) -invariant shapes. Then $C(X) \rtimes G$ is \mathcal{Z} -stable.*

Proof. Let $n \in \mathbb{N}$. Let Υ be a finite subset of the unit ball of $C(X)$, F a symmetric finite subset of G containing e , and $\varepsilon > 0$. We will show the existence of a complete positive contractive linear map $\varphi : M_n \rightarrow C(X) \rtimes G$ and a v as in Theorem 5.1 where the finite set Ω there is taken to be $\Upsilon \cup \{u_s : s \in F\}$. Since $C(X) \rtimes G$ is generated as a C^* -algebra by the unit ball of $C(X)$ and the unitaries u_s for $s \in G$, we will thereafter be able to conclude by Theorem 5.1 that $C(X) \rtimes G$ is \mathcal{Z} -stable.

Equip X with a compatible metric d . Choose an integer $Q > n^2/\varepsilon$.

Let $\kappa > 0$ be small enough so that the quantity $\gamma := 1 - (1 - \kappa)^{1/2}$ satisfies $\gamma < (1 - \gamma)/(2n)$. Take a $\beta > 0$ such that $\beta < (2n)^{-1}$ and $(1 - \beta)^2 \geq (1 - \kappa)^{1/2}$. We may also assume that β is small enough so that if T is a nonempty finite subset of G which is sufficiently invariant under left translation by F^{Q+1} and T' is a subset of T with $|T'| \geq (1 - n\beta)|T|$ then $|\bigcap_{s \in F^{Q+1}} s^{-1}T'| \geq (1 - \gamma)|T|$.

Choose an $L \in \mathbb{N}$ large enough so that $(1 - \beta/2)^L < \beta$. By amenability there exist finite subsets $e \in T_1 \subseteq T_2 \subseteq \dots \subseteq T_L$ of G such that $|\partial_{T_{l-1}}T_l| \leq (\beta/8)|T_l|$ for $l = 2, \dots, L$. By the previous paragraph, we may also assume that for each l the set T_l is sufficiently invariant under left translation by F^{Q+1} so that

$$(5.1) \quad \left| \bigcap_{s \in F^{Q+1}} s^{-1}T \right| \geq (1 - \gamma)|T|.$$

for all $T \subseteq T_l$ satisfying $|T| \geq (1 - n\beta)|T_l|$.

By uniform continuity there is a $\eta > 0$ such that $|f(x) - f(y)| < \varepsilon/(3n^2)$ for all $f \in \Upsilon \cup \Upsilon^2$ and all $x, y \in X$ satisfying $d(x, y) < \eta$. Again by uniform continuity there is an $\eta' > 0$ such that $d(tx, ty) < \eta$ for all $x, y \in X$ satisfying $d(x, y) < \eta'$ and all $t \in \bigcup_{l=1}^L T_l$. Fix a clopen partition $\{A_1, \dots, A_M\}$ of X whose members all have diameter less than η' .

Let E be a finite subset of G containing T_L and let $\delta > 0$ be such that $\delta \leq \beta/4$. We will further specify E and δ below. By hypothesis there is a collection $\{(V_k, S_k)\}_{k=1}^K$ of clopen towers such that the shapes S_1, \dots, S_K are (E, δ) -invariant and the sets $S_1 V_1, \dots, S_J V_J$ partition X . By a simple procedure we can construct, for each k , a clopen partition \mathcal{P}_k of V_k such that each level of every one of the towers (V, S_k) for $V \in \mathcal{P}_k$ is contained in one of the sets A_1, \dots, A_M . By replacing (V_k, S_k) with these thinner towers for each k , we may therefore assume that each level in every one of the towers $(V_1, S_1), \dots, (V_K, S_K)$ is contained in one of the sets A_1, \dots, A_M .

Let $1 \leq k \leq K$. Since S_k is $(T_L, \beta/4)$ -invariant, by Theorem 5.2 and our choice of the sets T_1, \dots, T_L we can find $C_{k,1}, \dots, C_{k,L} \subseteq S_k$ such that the collection $\{T_l c : l = 1, \dots, L, c \in C_{k,l}\}$ is β -disjoint and $(1 - \beta)$ -covers S_k . By β -disjointness, for every $l = 1, \dots, L$ and $c \in C_{k,l}$ we can find a $T_{k,l,c} \subseteq T_l$ satisfying $|T_{k,l,c}| \geq (1 - \beta)|T_l|$ so that the collection of sets $T_{k,l,c}$ for $l = 1, \dots, L$ and $c \in C_{k,l}$ is disjoint and has the same union as the sets $T_l c$ for $l = 1, \dots, L$ and $c \in C_{k,l}$, so that it $(1 - \beta)$ -covers S_k .

For each $l = 1, \dots, L$ and $m = 1, \dots, M$ write $C_{k,l,m}$ for the set of all $c \in C_{k,l}$ such that $cV_k \subseteq A_m$, and choose pairwise disjoint subsets $C_{k,l,m}^{(1)}, \dots, C_{k,l,m}^{(n)}$ of $C_{k,l,m}$ such that each has cardinality $\lfloor |C_{k,l,m}|/n \rfloor$. For each $i = 2, \dots, n$ choose a bijection

$$\Lambda_{k,i} : \bigsqcup_{l,m} C_{k,l,m}^{(1)} \rightarrow \bigsqcup_{l,m} C_{k,l,m}^{(i)}$$

which sends $C_{k,l,m}^{(1)}$ to $C_{k,l,m}^{(i)}$ for all l, m . Also, define $\Lambda_{k,1}$ to be the identity map from $\bigsqcup_{l,m} C_{k,l,m}^{(1)}$ to itself. Set $C'_{k,l,m} = \bigsqcup_{i=1}^n C_{k,l,m}^{(i)}$.

Let $1 \leq l \leq L$ and $c \in \bigsqcup_m C'_{k,l,m}$. Define the set $T'_{k,l,c} = \bigcap_{i=1}^n T_{k,l,\Lambda_{k,i}(c)}$, which satisfies

$$(5.2) \quad |T'_{k,l,c}| \geq (1 - n\beta)|T_l|$$

since each $T_{k,l,\Lambda_{k,i}(c)}$ has cardinality at least $(1 - \beta)|T_l|$. Set

$$B_{k,l,c,0} = T'_{k,l,c} \setminus F^{Q-1} B_{k,l,c,Q}, \quad B_{k,l,c,Q} = \bigcap_{s \in F^{Q+1}} sT'_{k,l,c}$$

and, for $q = 1, \dots, Q - 1$,

$$B_{k,l,c,q} = F^{Q-q} B_{k,l,c,Q} \setminus F^{Q-q-1} B_{k,l,c,Q}.$$

Then the sets $B_{k,l,c,0}, \dots, B_{k,l,c,Q}$ partition $T'_{k,l,c}$. For $s \in F$ we have

$$(5.3) \quad sB_{k,l,c,Q} \subseteq B_{k,l,c,Q-1} \cup B_{k,l,c,Q},$$

while for $q = 1, \dots, Q - 1$ we have

$$(5.4) \quad sB_{k,l,c,q} \subseteq B_{k,l,c,q-1} \cup B_{k,l,c,q} \cup B_{k,l,c,q+1},$$

for if we are given a $t \in B_{k,l,c,q}$ then $st \in F^{Q-q+1}B_{k,l,c,Q}$, while if $st \in F^{Q-q-2}B_{k,l,c,Q}$ then $t \in F^{Q-q-1}B_{k,l,c,Q}$ since F is symmetric, contradicting the membership of t in $B_{k,l,c,q}$. Also, from (5.1) and (5.2) we get

$$(5.5) \quad |B_{k,l,c,Q}| \geq (1 - \gamma)|T_l|.$$

For $i = 1, \dots, n$, $c \in \bigsqcup_m C_{k,l,m}^{(i)}$, and $q = 0, \dots, Q$ we set $B_{k,l,c,q} = B_{k,l,\lambda_{k,i}^{-1}(c),q}$.

Write $\Lambda_{k,i,j}$ for the composition $\Lambda_{k,i} \circ \Lambda_{k,j}^{-1}$. Define a linear map $\psi : M_n \rightarrow C(X) \rtimes G$ by declaring it on the standard matrix units $\{e_{ij}\}_{i,j=1}^n$ of M_n to be given by

$$\psi(e_{ij}) = \sum_{c \in C_{k,l,m}^{(j)}} \sum_{t \in T'_{k,l,c}} u_{t\Lambda_{k,i,j}(c)c^{-1}t^{-1}} \mathbf{1}_{tcV_k}$$

and extending linearly. Then $\varphi(e_{ij})^* = \varphi(e_{ji})$ for all i, j and the product $\psi(e_{ij})\psi(e_{i'j'})$ is 1 or 0 depending on whether $i = i'$, so that ψ is a $*$ -homomorphism.

For all k and l , all $1 \leq i, j \leq n$, and all $c \in \bigsqcup_m C_{k,l,m}^{(i)}$ we set

$$h_{k,l,c,i,j} = \sum_{q=1}^Q \sum_{t \in B_{k,l,c,q}} \frac{q}{Q} u_{t\Lambda_{k,i,j}(c)c^{-1}t^{-1}} \mathbf{1}_{tcV_k}$$

and put

$$h = \sum_{k,l,m} \sum_{i=1}^n \sum_{c \in C_{k,l,m}^{(i)}} h_{k,l,c,i,i}.$$

Then h is a norm-one function which commutes with the image of ψ , and so we can define an order-zero completely positive contractive linear map $\varphi : M_n \rightarrow C(X) \rtimes G$ by setting

$$\varphi(x) = h\psi(x).$$

Note that $\varphi(e_{ij}) = \sum_{k,l,m} \sum_{c \in C_{k,l,m}^{(j)}} h_{k,l,c,i,j}$.

We now verify condition (iii) in Theorem 5.1 for the elements of the set $\{u_s : s \in F\}$. Let $1 \leq i, j \leq n$. For all k and l , all $c \in \bigsqcup_m C_{k,l,m}^{(i)}$, and all $s \in F$ we have

$$\begin{aligned} u_s h_{k,l,c,i,j} u_s^{-1} - h_{k,l,c,i,j} &= \sum_{q=1}^Q \sum_{t \in B_{k,l,c,q}} \frac{q}{Q} u_{st\Lambda_{k,i,j}(c)c^{-1}(st)^{-1}} \mathbf{1}_{stcV_k} \\ &\quad - \sum_{q=1}^Q \sum_{t \in B_{k,l,c,q}} \frac{q}{Q} u_{t\Lambda_{k,i,j}(c)c^{-1}t^{-1}} \mathbf{1}_{tcV_k}, \end{aligned}$$

and so in view of (5.3) and (5.4) we obtain

$$\|u_s h_{k,l,c,i,j} u_s^{-1} - h_{k,l,c,i,j}\| \leq \frac{1}{Q} < \frac{\varepsilon}{n^2}.$$

Since each of the elements $a = u_s h_{k,l,c,i,j} u_s^{-1} - h_{k,l,c,i,j}$ is such that a^*a and aa^* are dominated by twice the indicator functions of $T'_{k,l,c}cV_k$ and $T'_{k,l,\Lambda_{k,i,j}(c)}\Lambda_{k,i,j}(c)V_k$, respectively,

and the sets $T'_{k,l,c}cV_k$ over all k and l and all $c \in \bigsqcup_m C'_{k,l,m}$ are pairwise disjoint, this yields

$$\|u_s \varphi(e_{ij})u_s^{-1} - \varphi(e_{ij})\| = \max_{k,l,m} \max_{c \in C_{k,l,m}^{(i)}} \|u_s h_{k,l,c,i,j} u_s^{-1} - h_{k,l,c,i,j}\| < \frac{\varepsilon}{n^2}$$

and hence, for every norm-one element $b = (b_{ij}) \in M_n$,

$$\begin{aligned} \|[u_s, \varphi(b)]\| &= \|u_s \varphi(b)u_s^{-1} - \varphi(b)\| \leq \sum_{i,j=1}^n |b_{ij}| \|u_s \varphi(e_{ij})u_s^{-1} - \varphi(e_{ij})\| \\ &< n^2 \cdot \frac{\varepsilon}{n^2} = \varepsilon. \end{aligned}$$

Next we verify condition (iii) in Theorem 5.1 for the functions in Υ . Let $1 \leq i, j \leq n$. Let $g \in \Upsilon \cup \Upsilon^2$. Let $1 \leq k \leq K$ and $1 \leq l \leq L$, and let $c \in C_{k,l,m}^{(j)}$. Then

$$(5.6) \quad h_{k,l,c,i,j}^* g h_{k,l,c,i,j} = \sum_{q=1}^Q \sum_{t \in B_{k,l,c,q}} \frac{q^2}{Q^2} (tc\Lambda_{k,i,j}(c)^{-1}t^{-1}g)\mathbf{1}_{tcV_k}$$

and

$$(5.7) \quad g h_{k,c,i,j}^* h_{k,c,i,j} = \sum_{q=1}^Q \sum_{t \in B_{k,l,c,q}} \frac{q^2}{Q^2} g \mathbf{1}_{tcV_k}$$

Now let $x \in V_k$. Since $\Lambda_{k,i,j}(c)x$ and cx both belong to A_m , we have $d(\Lambda_{k,i,j}(c)x, cx) < \eta'$. It follows that for every $t \in T_l$ we have $d(t\Lambda_{k,i,j}(c)x, tcx) < \eta$ by our choice of η' , so that $|g(t\Lambda_{k,i,j}(c)x) - g(tcx)| < \varepsilon/(3n^2)$ by our choice of η , in which case

$$\begin{aligned} \|(tc\Lambda_{k,i,j}(c)^{-1}t^{-1}g - g)\mathbf{1}_{tcV_k}\| &= \|c^{-1}t^{-1}((tc\Lambda_{k,i,j}(c)^{-1}t^{-1}g - g)\mathbf{1}_{tcV_k})\| \\ &= \|(\Lambda_{k,i,j}(c)^{-1}t^{-1}g - c^{-1}t^{-1}g)\mathbf{1}_{V_k}\| \\ &= \sup_{x \in V_k} |g(t\Lambda_{k,i,j}(c)x) - g(tcx)| \\ &< \frac{\varepsilon}{3n^2}. \end{aligned}$$

Using (5.6) and (5.7) this gives us

$$(5.8) \quad \begin{aligned} &\|h_{k,l,c,i,j}^* g h_{k,l,c,i,j} - g h_{k,l,c,i,j}^* h_{k,l,c,i,j}\| \\ &= \max_{q=1, \dots, Q} \max_{t \in B_{k,l,c,q}} \frac{q^2}{Q^2} \|(tc\Lambda_{k,i,j}(c)^{-1}t^{-1}g - g)\mathbf{1}_{tcV_k}\| < \frac{\varepsilon}{3n^2}. \end{aligned}$$

Set $w = \varphi(e_{ij})$ for brevity. Let $f \in \Upsilon$. Since the functions $h_{k,l,c,i,j}$ over all k, l , and $c \in C_{k,l,m}^{(i)}$ have pairwise disjoint supports, we infer from (5.8) that

$$\|w^* g w - g w^* w\| < \frac{\varepsilon}{3n^2}$$

for g equal to either f or f^2 . It follows that

$$\|w^* f^2 w - f w^* f w\| \leq \|w^* f^2 w - f^2 w^* w\| + \|f(f w^* w - w^* f w)\| < \frac{2\varepsilon}{3n^2}$$

and hence

$$\begin{aligned}
\|fw - wf\|^2 &= \|(fw - wf)^*(fw - wf)\| \\
&= \|w^*f^2w - fw^*fw + fw^*wf - w^*fwf\| \\
&\leq \|w^*f^2w - fw^*fw\| + \|(fw^*w - w^*fw)f\| \\
&< \frac{2\varepsilon}{3n^2} + \frac{\varepsilon}{3n^2} = \frac{\varepsilon}{n^2}.
\end{aligned}$$

Therefore for every norm-one element $b = (b_{ij}) \in M_n$ we have

$$\|[f, \varphi(b)]\| \leq \sum_{i,j=1}^n |b_{ij}| \|[f, \varphi(e_{ij})]\| < n^2 \cdot \frac{\varepsilon}{n^2} = \varepsilon.$$

Finally, we construct a $v \in C(X) \rtimes G$ that will satisfy conditions (i) and (ii) in Theorem 5.1. Let $1 \leq k \leq K$. For $i = 1, \dots, n$ define $S_{k,i} = \bigsqcup_{l,m} \bigsqcup_{c \in C_{k,l,m}^{(i)}} B_{k,l,c,Q}c$, which is the set of all $s \in S_k$ such that the function $\varphi(e_{ii})$ takes the value 1 on sV_k . Note that the sets $S_{k,1}, \dots, S_{k,n}$ all have the same cardinality. Set $S_{k,0} = S_k \setminus (S_{k,1} \cup \dots \cup S_{k,n})$. Since $|C_{k,l,m}^{(1)}| \geq \frac{1}{n}|C_{k,l,m}| - n$ for every l and m , choosing the sets T_1, \dots, T_n to be large enough (which we may do) and β to be small enough we can ensure that each of the sets $T'_{k,l,c}$ is large enough so that for every $i = 1, \dots, n$ we have

$$\left| \bigsqcup_{l,m} T'_{k,l,c} C_{k,l,m}^{(1)} \right| \geq \left(\frac{1}{n} - \gamma \right) \left| \bigsqcup_{l,m} T'_{k,l,c} C_{k,l,m} \right| \geq \left(\frac{1}{n} - \gamma \right) (1 - n\beta) |S_k|$$

with the second inequality coming from (5.2). Since for all l and all $c \in C_{k,l,m}^{(1)}$ we have

$$|B_{k,l,c,Q}| \geq (1 - \gamma) |T'_{k,l,c}|$$

it follows, putting $\lambda = (1 - \gamma)(1/n - \gamma)(1 - n\beta)$, that

$$(5.9) \quad |S_{k,1}| \geq (1 - \gamma) \left| \bigsqcup_{l,m} T'_{k,l,c} C_{k,l,m}^{(1)} \right| \geq \lambda |S_k| = \lambda (|S_{k,0}| + n|S_{k,1}|).$$

By taking γ and β small enough we can guarantee that $\lambda > 1/(n+1)$, in which case (5.9) implies that $|S_{k,0}| < |S_{k,1}|$. Hence there exists an injection $\theta_k : S_{k,0} \rightarrow S_{k,1}$. Define

$$z = \sum_{k=1}^K \sum_{s \in S_{k,0}} u_{\theta_k(s)s^{-1}} \mathbf{1}_{sV_k},$$

and note that z^*z is the indicator function of the set $\bigsqcup_{k=1}^K S_{k,0}V_k$ while zz^* is the indicator function of the set $\bigsqcup_{k=1}^K S_{k,1}V_k$. Now put $v = (1 - \varphi(1))^{1/2}z^*$. Then $vv^* = 1 - \varphi(1)$ while

$$v^*v = z(1 - \varphi(1))z^* \leq zz^* \leq \varphi(e_{11}).$$

as desired. \square

Combining Theorems 5.1 and 4.2 yields the following.

Theorem 5.4. *Let G be a countably infinite amenable group and X the Cantor set. Then the set of all actions in $\text{FrMin}(G, X)$ whose crossed product is \mathcal{Z} -stable is comeager, and in particular nonempty.*

Question 5.5. Do any of the crossed products in Theorem 5.4 have tracial state space with compact extreme boundary (from which we would be able to conclude finite nuclear dimension by [1] and hence classifiability)? For $G = \mathbb{Z}$ there is a comeager orbit in $\text{FrMin}(G, X)$ under the conjugation action of the homeomorphism group of X , and this generic action is the universal odometer, which is uniquely ergodic (so that the crossed product has a unique tracial state) [9]. However, already for \mathbb{Z}^2 nothing of this nature seems to be known. On the other hand, it is known that the crossed products of free minimal actions of finitely generated nilpotent groups on compact metrizable spaces of finite covering dimension have finite nuclear dimension, and in particular are \mathcal{Z} -stable [20].

REFERENCES

- [1] J. Bosa, N. Brown, Y. Sato, A. Tikuisis, S. White and W. Winter. Covering dimension of C^* -algebras and 2-coloured classification. To appear in *Mem. Amer. Math. Soc.*
- [2] T. Ceccherini-Silberstein, P. de la Harpe, and R. I. Grigorchuk. Amenability and paradoxical decompositions for pseudogroups and discrete metric spaces. (Russian) *Tr. Mat. Inst. Steklova* **224** (1999), 68–111; translation in *Proc. Steklov Inst. Math.* **224** (1999), 57–97.
- [3] T. Downarowicz, D. Huczek, and G. Zhang. Tilings of amenable groups. To appear in *J. Reine Angew. Math.*
- [4] G. Elliott, G. Gong, H. Lin, and Z. Niu. On the classification of simple amenable C^* -algebras with finite decomposition rank, II. arXiv:1507.03437.
- [5] G. Elek and G. Lippner, *Borel oracles. An analytic approach to constant time algorithms*, Proc. Amer. Math. Soc. 138 (2010), 2939–2947.
- [6] G. Gong, H. Lin, and Z. Niu. Classification of finite simple amenable \mathcal{Z} -stable C^* -algebras. arXiv:1501.00135.
- [7] E. Guentner, R. Willett, and G. Yu. Dynamic asymptotic dimension: relation to dynamics, topology, coarse geometry, and C^* -algebras. To appear in *Math. Ann.*
- [8] I. Hirshberg and J. Orovitz. Tracially \mathcal{Z} -absorbing C^* -algebras. *J. Funct. Anal.* **265** (2013), 765–785.
- [9] M. Hochman. Genericity in topological dynamics. *Ergodic Theory Dynam. Systems* **28** (2008), 125–165.
- [10] X. Jiang and H. Su. On a simple unital projectionless C^* -algebra. *Amer. J. Math.* **121** (1999), 359–413.
- [11] D. Kerr and H. Li. *Ergodic Theory: Independence and Dichotomies*. To appear in the Springer Monographs in Mathematics series.
- [12] A. Kechris, S. Solecki, and S. Todorcevic, *Borel chromatic numbers*, Adv. in Math. 141 (1999), 1–44.
- [13] R. Lyons and F. Nazarov. Perfect matchings as IID factors on non-amenable groups. *European J. Combin.* **32** (2011), 1115–1125.
- [14] H. Matui and Y. Sato. Strict comparison and \mathcal{Z} -absorption of nuclear C^* -algebras. *Acta Math.* **209** (2012), 179–196.
- [15] H. Matui and Y. Sato. Decomposition rank of UHF-absorbing C^* -algebras. *Duke Math. J.* **163** (2014), 2687–2708.
- [16] I. Namioka, *Følner’s conditions for amenable semi-groups*, Math. Scand. 15 (1964), 18–28.
- [17] D. S. Ornstein and B. Weiss. Entropy and isomorphism theorems for actions of amenable groups. *J. Analyse Math.* **48** (1987), 1–141.
- [18] M. Rørdam and W. Winter. The Jiang–Su algebra revisited. *J. Reine Angew. Math.* **642** (2010), 129–155.

- [19] G. Szabó. The Rokhlin dimension of topological \mathbb{Z}^m -actions. *Proc. Lond. Math. Soc. (3)* **110** (2015), 673–694.
- [20] G. Szabo, Ji. Wu, and J. Zacharias. Rokhlin dimension for actions of residually finite groups. Preprint, 2014. arXiv:1408.6096.
- [21] A. Tikuisis, S. White, and W. Winter. Quasidiagonality of nuclear C^* -algebras. *Ann. of Math. (2)* **185** (2017), 229–284.
- [22] W. Winter. Nuclear dimension and \mathcal{Z} -stability of pure C^* -algebras. *Invent. Math.* **187** (2012), 259–342.