

# THE CONE OF CURVES OF K3 SURFACES REVISITED

SÁNDOR J KOVÁCS

## 1. INTRODUCTION

The following theorem was proved in [Kov94] over the complex numbers. It turns out that the proof given there works with very small adjustments in arbitrary characteristic. The purpose of this note is to verify this, that is, to prove the following.

**Theorem 1.1.** *Let  $X$  be a K3 surface of Picard number at least three over an algebraically closed field of arbitrary characteristic. Then one of the following mutually exclusive conditions are satisfied:*

- (1.1.1)  *$X$  does not contain any curve of negative self-intersection*
- (1.1.2)  *$\overline{NE}(X) = \sum \mathbb{R}_+ \ell$  where the sum runs over all smooth rational curves on  $X$ .*

DEFINITIONS AND NOTATION 1.2. Let  $k$  be an algebraically closed field of arbitrary characteristic. Everything will be defined over  $k$ .

A K3 surface is a smooth projective surface  $X$  such that  $\omega_X \simeq \mathcal{O}_X$  and  $h^1(X, \mathcal{O}_X) = 0$ .

The closed cone of effective curves inside  $N_1(X)$  is denoted by  $\overline{NE}(X)$ . For more details about the construction and basic properties of this should consult [Kol96, II.4].

A class  $\xi \in N_1(X)$  is called *integral* if it can be represented by a divisor on  $X$ . It is called *effective* (respectively *ample*) if it is integral and can be represented by an effective (respectively ample) divisor. A class  $\xi \in \overline{NE}(X)$  is called *extremal* if it cannot be written as the sum of two incomparable classes in  $\overline{NE}(X)$ . A class of a smooth rational curve is called a *nodal* class. The set of all nodal classes is denoted by  $\mathcal{N}(X)$ .

Let  $h$  be an ample class and define

$$\mathcal{Q}(X) := \{\xi \in N_1(X) \mid \xi \cdot h > 0, \xi \cdot \xi = 0\}$$

Note that by the positivity condition this is just half of a quadric cone.

---

Supported in part by NSF Grant DMS-0856185, and the Craig McKibben and Sarah Merner Endowed Professorship in Mathematics at the University of Washington.

The convex hull of a set will be denoted by  $\text{Conv}$ .

An open subset of the boundary of the cone is called *circular* if the cone is not locally finitely generated at any point in the open set.

**ACKNOWLEDGMENT.** This paper owes its existence to Max Lieblich who suggested that perhaps the original statement is also true in arbitrary characteristic.

## 2. SIMPLE FACTS

Let us start with an easy, well-known consequence of Riemann-Roch theorem:

**Lemma 2.1.** *Let  $a \in N_1(X)$  be an integral class. Then  $a \cdot a$  is an even integer. If furthermore  $a \cdot a \geq -2$ , then either  $a$  or  $-a$  is effective.*

*Proof.* As  $a$  is integral, there exists a line bundle  $\mathcal{L}$  representing  $a$ . By Riemann-Roch

$$(2.1.1) \quad h^0(X, \mathcal{L}) - h^1(X, \mathcal{L}) + h^0(X, \mathcal{L}^\vee) = \frac{1}{2} a \cdot a + 2$$

and hence  $a \cdot a$  has to be even.

If  $a \cdot a \geq -2$ , then the right hand side of (2.1.1) is positive, so either  $h^0(X, \mathcal{L}) > 0$  or  $h^0(X, \mathcal{L}^\vee) > 0$ , so either  $a$  or  $-a$  is effective.  $\square$

**Corollary 2.2.**  *$\mathcal{Q}(X) \subset \overline{NE}(X)$  and for any  $e \in \mathcal{Q}(X)$  the hyperplane  $(e \cdot \underline{\phantom{x}} = 0) = \{\xi \in N_1(X) \mid e \cdot \xi = 0\}$  is a supporting hyperplane of  $\text{Conv}(\mathcal{Q}(X))$ .*

*Proof.* The first statement follows directly from (2.1) and since for any  $\xi \in \text{Conv}(\mathcal{Q}(X))$  (effective) irreducible class,  $\xi \cdot e \geq 0$ , this also implies the second statement.  $\square$

**Corollary 2.3.** *Let  $e, d \in \overline{NE}(X)$  such that  $e \cdot e = 0$  and  $d \cdot d > 0$ . Then  $d$  is in the interior of  $\overline{NE}(X)$  and  $e \cdot d > 0$ .*

*Proof.* Since  $d$  is in the interior of  $\mathcal{Q}(X)$  and  $\mathcal{Q}(X) \subset \overline{NE}(X)$ , it follows that  $d$  is in the interior of  $\overline{NE}(X)$ . Then since  $(e \cdot \underline{\phantom{x}} = 0)$  is a supporting hyperplane,  $d$  cannot be contained in it and hence  $e \cdot d > 0$ .  $\square$

Next we establish that nodal rays can only accumulate along the cone generated by  $\mathcal{Q}(X)$ .

**Lemma 2.4.** *Let  $\{\ell_n\} \subset \mathcal{N}(X)$  be an infinite sequence of nodal classes such that  $\mathbb{R}_+ \ell_n \xrightarrow[n \rightarrow +\infty]{} \mathbb{R}_+ \xi$  for some  $\xi \in \overline{NE}(X)$ . Then  $\xi \cdot \xi = 0$ .*

*Proof.* Let  $\ell \in \mathcal{N}(X)$ . Then for infinitely many  $n \in \mathbb{N}$ ,  $\ell_n \neq \ell$ , so  $\ell_n \cdot \ell \geq 0$  and hence  $\xi \cdot \ell \geq 0$ . Applying this with  $\ell = \ell_n$  yields that

$\xi \cdot \xi \geq 0$ . On the other hand since  $\ell_n \cdot \ell_n < 0$ , it follows that  $\xi \cdot \xi \leq 0$  as well and so we must have  $\xi \cdot \xi = 0$ .  $\square$

**Corollary 2.5.** *Let  $h \in \overline{NE}(X)$  be an ample class and  $\varepsilon > 0$  a real number. Define*

$$\mathcal{Q}_\varepsilon(X) := \{\xi \in N_1(X) \mid \xi \cdot h = 1, \xi \cdot \xi \geq -\varepsilon\}.$$

*Then the number of nodal classes not contained in  $\text{Conv}(\mathcal{Q}_\varepsilon(X))$  is finite.*

*Proof.* The set  $\{\xi \in \overline{NE}(X) \mid \xi \cdot h = 1\}$  is compact by Kleiman's criterion [KM98, 1.18] and hence any infinite set contained in it has an accumulation point. By (2.4) all accumulation points have to be contained in  $\mathcal{Q}_\varepsilon(X)$ , which implies the desired statement.  $\square$

**Corollary 2.6.** *Let  $\xi \in \overline{NE}(X)$  be an extremal class which is not a multiple of a nodal class. Then  $\xi \cdot \xi = 0$ .*

*Proof.* It follows from Riemann-Roch and [Kol96, II.4.14] that

$$(2.6.1) \quad \overline{NE}(X) = \sum_{\ell \in \mathcal{N}(X)} \mathbb{R}_+ \ell + \text{Conv}(\mathcal{Q}(X)),$$

in particular  $\xi \cdot \xi \leq 0$ , and then for any  $\varepsilon > 0$ ,

$$(2.6.2) \quad \overline{NE}(X) = \sum_{\substack{\ell \in \mathcal{N}(X) \\ \ell \notin \text{Conv}(\mathcal{Q}_\varepsilon(X))}} \mathbb{R}_+ \ell + (\overline{NE}(X) \cap \text{Conv}(\mathcal{Q}_\varepsilon(X))),$$

where the above sum is finite by (2.5).

Suppose that  $\xi \cdot \xi < 0$ . Without loss of generality we may assume that  $\xi \cdot h = 1$  and choose an  $\varepsilon > 0$  such that  $\xi \notin \text{Conv}(\mathcal{Q}_\varepsilon(X))$ . Then  $\xi$  is a multiple of a nodal class by (2.6.2).  $\square$

### 3. SUBCONES GENERATED BY TWO ELEMENTS

The following is a simple, but important computation.

**Lemma 3.1.** *Let  $e, d$  be effective classes such that  $e$  is indecomposable and  $d \cdot d > 0$ . Let  $L$  be the 2-dimensional linear subspace generated by  $e$  and  $d$  in  $N_1(X)$  and  $\mathcal{C} = L \cap \overline{NE}(X)$ . Then*

- (3.1.1) *if  $e \cdot e = 0$ , then there exists an  $f \in \mathcal{C}$  such that  $f$  is effective,  $e \cdot f > 0$ ,  $e$  and  $f$  are on opposite sides of  $d$ , and  $f \cdot f = 0$ , and*
- (3.1.2) *if  $e \cdot e = -2$ , then there exists an  $f \in \mathcal{C}$  such that  $f$  is effective,  $e \cdot f > 0$ ,  $e$  and  $f$  are on opposite sides of  $d$ , and  $f \cdot f = 0$  or  $f \cdot f = -2$ .*

*Proof.* Let  $A = d \cdot d$ ,  $B = e \cdot d$ , and  $C = e \cdot e$ . Set  $f = \alpha d - \beta e$ . Then

$$f \cdot f = A\alpha^2 - 2B\alpha\beta + C\beta^2.$$

If  $C = e \cdot e = 0$ , then  $B = e \cdot d > 0$  by (2.3) and hence the equation

$$0 = A\alpha^2 - 2B\alpha\beta = \alpha(A\alpha - 2B\beta)$$

has a positive integer solution,  $\alpha = 2B$ ,  $\beta = A$  such that the class  $f = \alpha d - \beta e$  has  $f \cdot f = 0$ . Since  $\alpha d = f + \beta e$ ,  $f$  and  $e$  lie on opposite sides of  $d$  and it also follows that  $f$  is effective by (2.1) and since  $e \cdot f > 0$ .

If  $C = e \cdot e = -2$ , then set  $x = B\alpha/2 + \beta$ ,  $y = \alpha/2$ , and  $N = 2A + B^2$ . Then

$$\begin{aligned} f \cdot f &= A\alpha^2 - 2B\alpha\beta - 2\beta^2 = \\ &= -2((B\alpha/2 + \beta)^2 - (2A + B^2)(\alpha/2)^2) = -2(x^2 - Ny^2). \end{aligned}$$

Now if  $N$  is a square, then as above there are two effective solutions for  $f \cdot f = 0$  and they are on opposite sides of  $d$  and hence one of them is on the side of  $d$  opposite to  $e$ .

If  $N$  is not a square then finding an  $f$  with  $f \cdot f = -2$  is equivalent to solving Pell's equation  $x^2 - Ny^2 = 1$  [IR90, 17.5.2]. One may choose a solution with both  $x, y > 0$  which again ensures that  $e$  and  $f$  are on opposite sides of  $d$  and that  $e \cdot f > 0$ . This completes the proof.  $\square$

**Corollary 3.2.** *If  $\varrho(X) = 2$ , let  $\overline{NE}(X) = \mathbb{R}_+\xi + \mathbb{R}_+\eta$ . Then one of the following mutually exclusive cases hold:*

- (3.2.1) *Neither  $\mathbb{R}_+\xi$  nor  $\mathbb{R}_+\eta$  contain any effective classes.*
- (3.2.2) *Both  $\mathbb{R}_+\xi$  and  $\mathbb{R}_+\eta$  contain an effective class of 0 or -2 self-intersection.*

*Proof.* If  $\mathcal{N}(X) = \emptyset$ , then the decomposition in (2.6.1) implies that  $\overline{NE}(X) = \text{Conv}(\mathcal{Q}(X))$ . If there exists an integral (equivalently, effective) class in  $\mathcal{Q}(X)$ , then by (3.1.1) we are in case (3.2.2). If there are no integral classes in  $\mathcal{Q}(X)$  then we are in case (3.2.1).

If  $\mathcal{N}(X) \neq \emptyset$ , then by the decomposition in (2.6.1) and (3.1.2) we are in case (3.2.2).  $\square$

#### 4. K3 SURFACES CONTAINING A SMOOTH RATIONAL CURVE

**Theorem 4.1.** *Let  $X$  be a K3 surface and  $\xi \in \overline{NE}(X)$  an extremal vector. Assume that  $\varrho(X) \geq 3$  and  $X$  contains a smooth rational curve. Then*

- (4.1.1)  *$\overline{NE}(X)$  has no circular part,*

(4.1.2) *There exists a sequence  $\ell_n \in \mathcal{N}(X)$  such that  $\mathbb{R}_+ \ell_n \xrightarrow{n \rightarrow +\infty} \mathbb{R}_+ \xi$ ,*

*and*

$$(4.1.3) \quad \overline{NE}(X) = \overline{\sum_{\ell \in \mathcal{N}(X)} \mathbb{R}_+ \ell}.$$

*Proof.* Let  $\ell$  denote the class of a smooth rational curve, guaranteed by the assumption and let  $\sigma : X \rightarrow X'$  be the morphism contracting  $\ell$  to a point. Recall that  $X'$  is still projective and let  $h_1$  and  $h_2$  be two linearly independent ample classes on  $X'$ . Let  $d_i = \sigma^* h_i$  for  $i = 1, 2$  and observe that  $d_1$  and  $d_2$  are effective classes on  $X$  such that  $\ell \cdot d_1 = \ell \cdot d_2 = 0$  and  $\ell, d_1, d_2$  are linearly independent.

Suppose there exists  $U \subset \partial \overline{NE}(X)$  a non-empty open subset of  $\partial \overline{NE}(X)$  such that  $\mathbb{R}_+ U$  is a circular part (i.e., nowhere locally finitely generated) of  $\overline{NE}(X)$ . By (2.4) it follows that  $U \subseteq \mathcal{Q}(X)$  and hence in a neighbourhood of  $\mathbb{R}_+ U$  every effective class has non-negative self-intersection.

Let  $h$  be an arbitrary ample class and observe that by (3.1) the 2-dimensional linear subspace generated by  $h$  and  $\ell$  contains and  $f$  on the side of  $h$  opposite to  $\ell$  there is an effective class  $f$  with either  $f \cdot f = 0$  or  $f \cdot f = -2$ . We may repeat the same procedure with  $\ell$  replaced by  $f$  and  $h$  replaced by another ample class and find that these classes are all over near the boundary of  $\overline{NE}(X)$ . In particular, we can find an ample class  $h \in \overline{NE}(X)$  and an effective class  $f$  with either  $f \cdot f = 0$  or  $f \cdot f = -2$  such that the 2-dimensional linear subspace generated by  $h$  and  $f$  intersects  $U$  non-trivially. Then applying (3.1) again and combining it by the observation above we obtain that there exists an effective class  $e \in U$  such that  $e \cdot e = 0$ .

Next let  $d$  one of  $d_1$  and  $d_2$  such that  $e, d, \ell$  are linearly independent. Let  $A = d \cdot d$ ,  $B = e \cdot d$ , and  $C = e \cdot \ell$ . Recall that by choice of  $d$  we have  $d \cdot \ell = 0$ .

**CLAIM 4.1.4.**  $2B^2 \neq AC^2$

*Proof.* Suppose  $2B^2 = AC^2$  and let  $f = ACe - BCd + B^2\ell$ . Then  $f \cdot f = B^2(AC^2 - 2B^2) = 0$  and  $f \cdot e = 0$ . Then by (2.1)  $f$  or  $-f$  is effective. However, since  $e \in U$  which is a circular part of  $\partial \overline{NE}(X)$ , the only effective classes contained in the hyperplane ( $e \cdot \_ = 0$ ) are multiples of  $e$ . This implies that then  $BC = B^2 = 0$ , so  $B = 0$ . Applying the same argument for  $d$ , it would follow that  $d$  is a multiple of  $e$  which is impossible by the choice of  $d$  and  $e$ . Therefore we reached a contradiction and hence the claim is proven.  $\square$

Next let  $n \in \mathbb{N}$  and define

$$\pm d_n = (2(2B^2 - AC^2)Cn^2 - 4Bn)e + (2C^2n)d + (1 - 2BCn)\ell.$$

Then  $d_n \cdot d_n = -2$  and by (2.1) either  $d_n$  or  $-d_n$  is effective. Choose  $d_n$  to be effective. By (4.1.4)  $2B^2 - AC^2 \neq 0$  and hence  $\mathbb{R}_+d_n \rightarrow \mathbb{R}_+e$ , but this contradicts the observation that in a neighbourhood of  $U$  every effective class has non-negative self-intersection. Therefore (4.1.1) is proven.

Now let  $\xi \in \overline{NE}(X)$  extremal. If no multiple of  $\xi$  is in  $\mathcal{N}(X)$ , then  $\xi \in sQ(X)$  by (2.6). If  $\xi$  were not contained in the closure of the convex cone generated by  $\mathcal{N}(X)$ , then by [Kov94, 2.6]  $\overline{NE}(X)$  would have a circular part, so (4.1.2) follows from (4.1.1).

Finally, since every class in  $\overline{NE}(X)$  may be written as a sum of finitely many extremal classes (4.1.3) follows from (4.1.2).  $\square$

**Corollary 4.2.** *Let  $X$  be a K3 surface of Picard number at least three over an algebraically closed field of arbitrary characteristic. Then one of the following mutually exclusive conditions are satisfied:*

$$(4.2.1) \quad \overline{NE}(X) = \overline{\text{Conv}(\mathcal{Q}(X))}, \text{ or}$$

$$(4.2.2) \quad \overline{NE}(X) = \overline{\sum_{\ell \in \mathcal{N}(X)} \mathbb{R}_+ \ell}.$$

*Proof.* If  $X$  does not contain any curve of negative self-intersection, then (4.2.1) follows from (2.6.1). Otherwise (4.2.2) follows from (4.1).  $\square$

**REMARK 4.3.** Clearly (4.2) is equivalent to (1.1).

## REFERENCES

- [IR90] K. IRELAND AND M. ROSEN: *A classical introduction to modern number theory*, second ed., Graduate Texts in Mathematics, vol. 84, Springer-Verlag, New York, 1990. [1070716 \(92e:11001\)](#)
- [Kol96] J. KOLLÁR: *Rational curves on algebraic varieties*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, vol. 32, Springer-Verlag, Berlin, 1996. [MR1440180 \(98c:14001\)](#)
- [KM98] J. KOLLÁR AND S. MORI: *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. [MR1658959 \(2000b:14018\)](#)
- [Kov94] S. J. KOVÁCS: *The cone of curves of a K3 surface*, *Math. Ann.* **300** (1994), no. 4, 681–691. [MR1314742 \(96a:14044\)](#)

UNIVERSITY OF WASHINGTON, DEPARTMENT OF MATHEMATICS, BOX 354350,  
SEATTLE, WA 98195-4350, USA

*E-mail address:* [skovacs@uw.edu](mailto:skovacs@uw.edu)

*URL:* <http://www.math.washington.edu/~kovacs>